



GOVERNMENT OF THE DISTRICT OF COLUMBIA
OFFICE OF THE CHIEF MEDICAL EXAMINER

401 E Street, SW – 6th Floor
Washington, DC 20024


Opioid-related Fatal Overdoses: January 1, 2014 to May 31, 2016¹

Report Date: August 10, 2016

The DC Office of the Chief Medical Examiner (OCME) investigated a total of **269²** deaths due to use of opioids from January 1, 2014 through April 30, 2016, **83** deaths in CY 2014 and **114** in CY 2015 and **72** deaths to date in CY 2016 respectively. This report examines the presence of opioids (*heroin, fentanyl, fentanyl analogs, morphine, prescription opioids and the general classification of opiates*) in the deaths observed at the OCME. The tables and graphs below present the decedent information by trends, demographics, and jurisdiction of residence.


Incidence of Opioids by Year

Each drug is counted independently in fatalities involving more than one of these drugs and ranged from 1 to 5 opioids identified per death. Therefore, there were a total of **120** opioids³ found in the 83 deaths in 2014, **160** opioids found in the 114 deaths in 2015 and **117** opioids found in the 72 deaths in 2016. As depicted in the graph to the right, the total number of opioids that caused a death increased from 2014 to 2015. If current trends persist, the drug count in 2016 will exceed the total number of drugs observed in 2015.



Trends in Deaths due to Opioid Use

Examining the monthly trends in deaths caused by opioid drug use, more fatalities occurred in the month of May of 2016 than any other month over the past five years. Overall, there was a 37% increase in fatal overdoses due to opioid use from 2014 (n=83) to 2015 (n=114). The current trends in 2016 surpass the prevalence of opioid-related overdoses during the same time period in that given year.



¹ Data for 2016 is inconclusive and subject to change due to cases where cause and manner of death is “Pending Further Investigation”

² The data presented in this report includes 10 cases with a Manner of Death other than Accidental Intoxication- three cases in 2014, one case in 2015, and one case in 2016 in which the Manner of Death was Undetermined but the Cause of Death was due to opioid drug use. Additionally there were 2 cases with Manner of Death of Suicide in 2014, one case in 2015 and two cases in 2016.

³ Morphine and fentanyl can both be prescribed. However, for the sake of this report, they are included under the illicit opioids



Increase in Cases since September

Seventy-five percent of deaths involving acetyl fentanyl were found among decedents between the months of September through December. Of the nine cases of acetyl fentanyl since September 2015, there were 3 decedents that were residents of Ward 4, 2 decedents that were undomiciled and the remaining four decedents were spread equally in Wards 5, 6, 8, or were jurisdiction of residence was unknown. In 2016, **57%** of the cases in March through May had either fentanyl or fentanyl analog (fentanyl, furanyl-fentanyl or despropionyl-fentanyl) on board.



Demographics¹

Race/Ethnicity

Overall, **203** or **75%** of all deaths due to opioid use were among Blacks. Moreover, this trend is remains when examining the data across years.




Gender

Fatal overdoses due to opioid drug use were most common among males than females across years.





Age

Approximately, **78%** of all overdoses due to opioid drug use happen among adults between the ages of 40-69 years old. Overall, deaths due to opioid use were most prevalent among people ages 50 to 59 (41%). There were no deaths from the use of opioids among individuals younger than 16 and older than 89.






Jurisdiction of Residence¹

Overall, opioid-related fatal overdoses were most prevalent in Ward 8. However, there are variations across study years. For example; in opioid-related fatal overdoses were most prevalent in wards 7 and 8 in 2015 compared to Wards 5 and 7 in 2014. The largest increase from 2014 to 2015 was observed in Ward 8, which increased by over 250%. In 2016 to date, there are more opioid-related overdoses in Wards 7 and 8.



The graphs to the right depict the total number of heroin, morphine, fentanyl, and fentanyl analogs that contributed to death by Ward of Residence. Each drug is counted independently in fatalities involving more than one of these drugs. The total number of opioids found in fatal overdoses increased between 2014 and 2015. Overall, in the District there were a total of **64** counts of heroin, morphine, and fentanyl that contributed to fatal overdoses in 2014. However, in 2015, the number of opioids that contributed to a fatal overdose in the District increased to a total of **100** opioids (heroin, morphine, fentanyl, and acetyl fentanyl). Fentanyl is a synthetic and short acting opioid analgesic that has been approved for managing acute or chronic pain. Fentanyl and fentanyl analogs are highly potent synthetic opioids, which are more potent than morphine and heroin. According to the CDC, acetyl fentanyl is a synthetic opioid equivalent fentanyl that is up to five times more potent than heroin.⁴ There were zero cases of acetyl fentanyl in 2014. In contrast, nine of the twelve cases of acetyl fentanyl found in 2015 were among decedents that were residence of the District. To




⁴ <http://www.emergency.cdc.gov/HAN/han00350.asp>



date in 2016, there were a total of 63 opioids (heroin, morphine, fentanyl, furanyl-fentanyl and despropionyl-fentanyl). Although, there are zero cases of acetyl fentanyl in 2016, new fentanyl analogs have emerged that are equally as potent.


Prescription Opioids

There were 90 prescription opioids found in the 244 drug overdoses between 2014 and 2015. Fatal overdoses involving prescription opioids were more prevalent in 2014 (n=45) than in 2015 (n=29). This represents a 36% decrease in prescription drugs that contributed to drug overdoses. Overall, the most commonly detected prescription Opioid was methadone. However, there are differences when you examine the data by year. In 2014, methadone was the most commonly found prescription opioid in drug overdoses. However, in 2015, oxycodone was the most prevalent prescription opioid identified. Currently in 2016, there are 22 incidences of prescription opioids that contributed to a fatal overdose, which suggests an increasing trend from 2015.



Map of Opioid related Deaths by Jurisdiction of Residence

The map below displays the drug overdoses due to opioid use by jurisdiction of residence. The jurisdiction of residence is not known for all decedents reported to and investigated by OCME. This map includes all decedents that died in the District of Columbia where the address was known, which is approximately 77% of all opioid related drug deaths between 2014 and 2015. There was not any Ward in 2014 and 2015 that was exempt from having a resident experience a fatal opioid overdose.



Office of the Chief Medical Examiner
401 E. Street SW, Washington, DC 20024

Rev. 8/10/16

Map of Opioid related Deaths by Location of Injury

The map below displays the drug overdoses due to Opioid use by location of injury. The location of injury is not known for all decedents reported and investigated by OCME. This map includes all decedents where the location of injury is known, which is approximately 70% all Opioid related deaths (n=138). A fatal opioid overdose occurred in every Ward in 2014 and 2015.

